

Press Release

ENHERTU[®] Approved in the U.S. as First HER2 Directed Therapy for Patients with HER2 Low or HER2 Ultralow Metastatic Breast Cancer Following Disease Progression After One or More Endocrine Therapies

- Based on DESTINY-Breast06 phase 3 trial results which showed ENHERTU demonstrated superiority versus chemotherapy with a median progression-free survival of more than one year
- Approval brings Daiichi Sankyo and AstraZeneca's ENHERTU to an earlier HR positive treatment setting and broadens the patient population eligible for treatment with a HER2 directed therapy to those with HER2 ultralow disease

Tokyo and Basking Ridge, NJ – (January 27, 2025) – Daiichi Sankyo (TSE: 4568) and AstraZeneca's (LSE/STO/Nasdaq: AZN) ENHERTU[®] (fam-trastuzumab deruxtecan-nxki) has been approved in the U.S. for the treatment of adult patients with unresectable or metastatic hormone receptor (HR) positive, HER2 low (IHC 1+ or IHC 2+/ISH-) or HER2 ultralow (IHC 0 with membrane staining) breast cancer, as determined by an FDA-approved test, that has progressed on one or more endocrine therapies in the metastatic setting.

ENHERTU is a specifically engineered HER2 directed DXd antibody drug conjugate (ADC) discovered by Daiichi Sankyo and being jointly developed and commercialized by Daiichi Sankyo and AstraZeneca.

This approval, which is based on results from the [DESTINY-Breast06](#) phase 3 trial [presented](#) at the 2024 American Society of Clinical Oncology (#ASCO24) Annual Meeting and published in [The New England Journal of Medicine](#), follows [Priority Review](#) and [Breakthrough Therapy Designation](#) by the FDA for ENHERTU in this indication.

In the DESTINY-Breast06 trial, ENHERTU demonstrated a 36% reduction in the risk of disease progression or death versus chemotherapy in the overall trial population of patients with chemotherapy-naïve HR positive, HER2 low or HER2 ultralow metastatic breast cancer (n=866) (hazard ratio [HR] 0.64; 95% confidence interval [CI]: 0.54-0.76; p<0.0001). A median progression-free survival (PFS) of 13.2 months (95% CI: 12.0-15.2) was seen in patients treated with ENHERTU compared to 8.1 months (95% CI: 7.0-9.0) in patients treated with chemotherapy.

The confirmed objective response rate (ORR) in the overall trial population was 62.6% (95% CI: 57.6-67.4) in the ENHERTU arm versus 34.4% (95% CI: 29.7-39.4) in the chemotherapy arm as assessed by blinded independent central review (BICR). There were 10 (2.5%) complete responses (CRs) and 236 (60.1%) partial responses (PRs) seen in the ENHERTU arm compared to zero (0%) CRs and 134 (34.4%) PRs in the chemotherapy arm. The median duration of response (DOR) was 14.3 months in the ENHERTU arm (95% CI: 12.5-15.9) versus 8.6 months in the chemotherapy arm (95% CI: 6.9-11.5) as assessed by BICR.

In the HER2 low population (n=713) (HR 0.62; 95% CI: 0.52-0.75; p<0.0001), median PFS was 13.2 months (95% CI: 11.4-15.2) in the ENHERTU arm compared to 8.1 months (95% CI: 7.0-9.0) in the chemotherapy arm. Confirmed ORR was 62.0% (95% CI: 56.5-67.3) versus 35.2% (95% CI: 30.0-40.7) in the chemotherapy arm. There were 9 (2.8%) CRs and 193 (59.2%) PRs seen in the ENHERTU arm compared to zero (0%) CRs and 114 (35.2%) PRs in the chemotherapy arm. Median DOR was 14.1 months (95% CI: 11.9-15.9) versus 8.6 months (95% CI: 6.7-11.3) in the chemotherapy arm.

An exploratory analysis of the HER2 ultralow population (n=153) (HR 0.76; 95% CI: 0.49-1.17) showed the clinically meaningful improvement in PFS was consistent between patients with HER2 low and HER2 ultralow expression, with 15.1 months (95% CI: 10.0-17.3) reported in the ENHERTU arm and 8.3 months (95% CI: 5.8-15.2) in the chemotherapy arm. Confirmed ORR was 65.7% (95% CI: 53.1-76.8) in the ENHERTU arm and 30.8% (95% CI: 19.9-43.4) in the chemotherapy arm. Median DOR was 14.3 months (95% CI: 11.8-not estimable [NE]) versus 14.1 months (95% CI: 5.9-NE) in the chemotherapy arm.

“Endocrine therapy is typically used in the initial treatment of HR positive metastatic breast cancer and following progression, subsequent chemotherapy is associated with poor outcomes,” said Aditya Bardia, MD, MPH, Program Director of Breast Oncology and Director of Translational Research Integration, UCLA Health Jonsson Comprehensive Cancer Center and Investigator in the DESTINY-Breast06 trial. “With a median progression-free survival exceeding one year and a response rate of more than 60 percent, trastuzumab deruxtecan offers a potential new standard of care for patients with hormone receptor positive, HER2 low or HER2 ultralow metastatic breast cancer following endocrine therapy.”

“We are excited to see more treatment options for these patients which enable more personalized care,” said Krissa Smith, Vice President, Education, Susan G. Komen. “It is critical for patients to understand the HER2 status of their metastatic breast cancer to help them make informed treatment decisions. Patients with tumors that are HER2 low or HER2 ultralow now have more options to consider with their healthcare team.”

ENHERTU is approved with Boxed WARNINGS for interstitial lung disease (ILD)/pneumonitis and Embryo-Fetal toxicity. The safety of ENHERTU was evaluated in 434 patients with unresectable or metastatic HER2 low (IHC 1+ or IHC 2+/ISH-) or HER2 ultralow (IHC 0 with membrane staining) breast

cancer who received ENHERTU (5.4 mg/kg) in DESTINY-Breast06. The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased white blood cell count, decreased neutrophil count, nausea, decreased hemoglobin, decreased lymphocyte count, fatigue, decreased platelet count, alopecia, increased alanine aminotransferase, increased blood alkaline phosphatase, increased aspartate aminotransferase, decreased blood potassium, diarrhea, vomiting, constipation, decreased appetite, COVID-19 and musculoskeletal pain. Serious adverse reactions occurred in 20% of patients receiving ENHERTU. Serious adverse reactions in $>1\%$ of patients who received ENHERTU were ILD/pneumonitis, COVID-19, febrile neutropenia and hypokalemia. Fatalities due to adverse reactions occurred in 2.8% of patients.

“ENHERTU continues to redefine the classification and treatment of HR positive metastatic breast cancer with important new data across the spectrum of HER2 expression,” said Ken Keller, Global Head of Oncology Business, and President and CEO, Daiichi Sankyo, Inc. “Today’s approval underscores our ongoing commitment to realizing the full potential of this innovative antibody drug conjugate and represents another paradigm shift in how certain breast cancers can be treated.”

“Building on the practice-changing previous approvals for ENHERTU, this new approval brings this important medicine to an earlier treatment setting and a broader patient population with HER2 expressing metastatic breast cancer,” said Dave Fredrickson, Executive Vice President, Oncology Hematology Business Unit, AstraZeneca. “The approval also highlights the importance of testing metastatic breast cancer tumors for detectable staining with a standard IHC test to identify those who may be eligible for treatment with ENHERTU following endocrine therapy.”

Daiichi Sankyo and AstraZeneca are committed to ensuring that patients in the U.S. who are prescribed ENHERTU can access the medication and receive necessary financial support. Provider and patient support, reimbursement and distribution for ENHERTU in the U.S. will be accessible by visiting www.ENHERTU4U.com or calling 1-833-ENHERTU (1-833-364-3788).

Please visit www.ENHERTU.com for full [Prescribing Information](#), including Boxed WARNINGS, and [Medication Guide](#).

Financial Considerations

Following this approval in the U.S., an amount of \$175 million is due from AstraZeneca to Daiichi Sankyo as a milestone payment for the HER2 low and HER2 ultralow chemotherapy-naïve breast cancer indication. Sales of ENHERTU in the U.S. are recognized by Daiichi Sankyo. For further details on the financial arrangements, please consult the collaboration agreement from [March 2019](#).

About DESTINY-Breast06

DESTINY-Breast06 is a global, randomized, open-label, phase 3 trial evaluating the efficacy and safety of ENHERTU (5.4 mg/kg) versus investigator's choice of chemotherapy (capecitabine, paclitaxel or nab paclitaxel) in patients with HR positive, HER2 low (IHC 1+ or IHC 2+/ISH-) or HER2 ultralow (defined as IHC 0 with membrane staining) advanced or metastatic breast cancer. Patients in the trial had no prior chemotherapy for advanced or metastatic disease and received at least two lines of prior endocrine therapy in the metastatic setting. Patients also were eligible if they had received one prior line of endocrine therapy combined with a CDK4/6 inhibitor in the metastatic setting and experienced disease progression within six months of starting first-line treatment or received endocrine therapy as an adjuvant treatment and experienced disease recurrence within 24 months.

HER2 IHC status was confirmed by a central laboratory and determined based on the most recent evaluable HER2 IHC sample prior to randomization. In tumor samples from patients screened for trial eligibility, nearly two-thirds of tumors previously assessed as IHC 0 at a local laboratory were re-classified as HER2 low or HER2 ultralow upon central analysis of the archival tumor sample. It was also observed that approximately 85% to 90% patients with HR positive, HER2 negative metastatic breast cancer may have actionable levels of HER2 expression.

The primary endpoint of DESTINY-Breast06 is PFS in the HR positive, HER2 low patient population as measured by BICR. Key secondary endpoints include PFS by BICR in the overall trial population (HER2 low and HER2 ultralow), OS in patients in the HER2 low patient population and OS in the overall trial population. Other secondary endpoints include ORR, DOR, time to first subsequent treatment or death, time to second subsequent treatment or death and safety. Analysis of the HER2 ultralow subgroup was not powered to demonstrate statistical significance.

DESTINY-Breast06 enrolled 866 patients (n=713 for HER2 low and n=153 for HER2 ultralow) in Asia, Europe, North America, Oceania and South America. For more information about the trial, visit [ClinicalTrials.gov](https://clinicaltrials.gov).

About Breast Cancer and HER2 Expression

Breast cancer is the second most common cancer and one of the leading causes of cancer-related deaths worldwide.¹ More than two million breast cancer cases were diagnosed in 2022 with more than 665,000 deaths globally.¹ In the U.S., more than 300,000 cases of breast cancer are diagnosed annually.² While survival rates are high for those diagnosed with early breast cancer, only about 30% of patients diagnosed with or who progress to metastatic disease are expected to live five years following diagnosis.³

HR positive, HER2 negative is the most common breast cancer subtype, accounting for approximately 70% of all breast cancers.³ HER2 is a tyrosine kinase receptor growth-promoting protein expressed on the surface

of many types of tumors, including breast cancer.⁴ Patients with high levels of HER2 expression (IHC 3+ or IHC2+/ISH+) are classified as HER2 positive and treated with HER2 targeted therapies, representing approximately 15% to 20% of all breast cancers.⁵ Historically, tumors that were not classified as HER2 positive were classified as HER2 negative, despite the fact that many of these tumors still carry some level of HER2 expression.⁶

Endocrine therapy is widely given consecutively in the early lines of treatment for HR positive metastatic breast cancer. However, after initial therapy, further efficacy with additional endocrine treatment is often limited.⁷ The current standard of care following endocrine therapy is chemotherapy, which is associated with poor response rates and outcomes.^{7,8,9,10}

Prior to the approvals of ENHERTU in HER2 low and HER2 ultralow metastatic breast cancer based on the [DESTINY-Breast04](#) and [DESTINY-Breast06](#) trials, there were no targeted therapies approved specifically for these patient populations.^{11,12}

About ENHERTU

ENHERTU (trastuzumab deruxtecan; fam-trastuzumab deruxtecan-nxki in the U.S. only) is a HER2 directed ADC. Designed using Daiichi Sankyo's proprietary DXd ADC Technology, ENHERTU is the lead ADC in the oncology portfolio of Daiichi Sankyo and the most advanced program in AstraZeneca's ADC scientific platform. ENHERTU consists of a HER2 monoclonal antibody attached to a number of topoisomerase I inhibitor payloads (an exatecan derivative, DXd) via tetrapeptide-based cleavable linkers.

ENHERTU (5.4 mg/kg) is approved in more than 75 countries worldwide for the treatment of adult patients with unresectable or metastatic HER2 positive (immunohistochemistry [IHC] 3+ or in-situ hybridization (ISH)+) breast cancer who have received a prior anti-HER2-based regimen, either in the metastatic setting or in the neoadjuvant or adjuvant setting, and have developed disease recurrence during or within six months of completing therapy based on the results from the [DESTINY-Breast03](#) trial.

ENHERTU (5.4 mg/kg) is approved in more than 75 countries worldwide for the treatment of adult patients with unresectable or metastatic HER2 low (IHC 1+ or IHC 2+/ISH-) breast cancer who have received a prior systemic therapy in the metastatic setting or developed disease recurrence during or within six months of completing adjuvant chemotherapy based on the results from the [DESTINY-Breast04](#) trial.

ENHERTU (5.4 mg/kg) is approved in the U.S. for the treatment of adult patients with unresectable or metastatic hormone receptor (HR) positive, HER2 low (IHC 1+ or IHC 2+/ISH-) or HER2 ultralow (IHC 0 with membrane staining) breast cancer, as determined by an FDA-approved test, that has progressed on one or more endocrine therapies in the metastatic setting based on the results from the [DESTINY-Breast06](#) trial.

ENHERTU (5.4 mg/kg) is approved in more than 50 countries worldwide for the treatment of adult patients with unresectable or metastatic NSCLC whose tumors have activating *HER2 (ERBB2)* mutations, as detected by a locally or regionally approved test, and who have received a prior systemic therapy based on the results from the [DESTINY-Lung02](#) and/or [DESTINY-Lung05](#) trials. Continued approval in China and the U.S. for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

ENHERTU (6.4 mg/kg) is approved in more than 65 countries worldwide for the treatment of adult patients with locally advanced or metastatic HER2 positive (IHC 3+ or IHC 2+/ISH+) gastric or gastroesophageal junction (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen based on the results from the [DESTINY-Gastric01](#), [DESTINY-Gastric02](#) and/or [DESTINY-Gastric06](#) trials. Continued approval in China for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

ENHERTU (5.4 mg/kg) is approved in Brazil, Israel, Russia and the U.S. for the treatment of adult patients with unresectable or metastatic HER2 positive (IHC 3+) solid tumors who have received prior systemic treatment and have no satisfactory alternative treatment options based on efficacy results from the [DESTINY-PanTumor02](#), [DESTINY-Lung01](#) and [DESTINY-CRC02](#) trials. Continued approval for this indication in the U.S. may be contingent upon verification and description of clinical benefit in a confirmatory trial.

About the ENHERTU Clinical Development Program

A comprehensive global clinical development program is underway evaluating the efficacy and safety of ENHERTU monotherapy across multiple HER2 targetable cancers. Trials in combination with other anticancer treatments, such as immunotherapy, also are underway.

About the Daiichi Sankyo and AstraZeneca Collaboration

Daiichi Sankyo and AstraZeneca entered into a global collaboration to jointly develop and commercialize ENHERTU in [March 2019](#) and DATROWAY® in [July 2020](#), except in Japan where Daiichi Sankyo maintains exclusive rights for each ADC. Daiichi Sankyo is responsible for the manufacturing and supply of ENHERTU and DATROWAY.

About the ADC Portfolio of Daiichi Sankyo

The Daiichi Sankyo ADC portfolio consists of seven ADCs in clinical development crafted from two distinct ADC technology platforms discovered in-house by Daiichi Sankyo.

The ADC platform furthest in clinical development is Daiichi Sankyo's DXd ADC Technology where each ADC consists of a monoclonal antibody attached to a number of topoisomerase I inhibitor payloads (an exatecan derivative, DXd) via tetrapeptide-based cleavable linkers. The DXd ADC portfolio currently consists of ENHERTU, a HER2 directed ADC, and DATROWAY, a TROP2 directed ADC, which are being jointly developed and commercialized globally with AstraZeneca. Patritumab deruxtecan (HER3-DXd), a HER3 directed ADC, ifinatamab deruxtecan (I-DXd), a B7-H3 directed ADC, and raludotatug deruxtecan (R-DXd), a CDH6 directed ADC, are being jointly developed and commercialized globally with Merck & Co., Inc, Rahway, NJ, USA. DS-3939, a TA-MUC1 directed ADC, is being developed by Daiichi Sankyo.

The second Daiichi Sankyo ADC platform consists of a monoclonal antibody attached to a modified pyrrolobenzodiazepine (PBD) payload. DS-9606, a CLDN6 directed PBD ADC, is the first of several planned ADCs in clinical development utilizing this platform.

Ifinatamab deruxtecan, patritumab deruxtecan, raludotatug deruxtecan, DS-3939 and DS-9606 are investigational medicines that have not been approved for any indication in any country. Safety and efficacy have not been established.

ENHERTU U.S. Important Safety Information

Indications

ENHERTU is a HER2-directed antibody and topoisomerase inhibitor conjugate indicated for the treatment of adult patients with:

- Unresectable or metastatic HER2-positive (IHC 3+ or ISH positive) breast cancer who have received a prior anti-HER2-based regimen either:
 - In the metastatic setting, or
 - In the neoadjuvant or adjuvant setting and have developed disease recurrence during or within six months of completing therapy
- Unresectable or metastatic:
 - Hormone receptor (HR)-positive, HER2-low (IHC 1+ or IHC 2+/ISH-) or HER2-ultralow (IHC 0 with membrane staining) breast cancer, as determined by an FDA-approved test, that has progressed on one or more endocrine therapies in the metastatic setting
 - HER2-low (IHC 1+ or IHC 2+/ISH-) breast cancer, as determined by an FDA-approved test, who have received a prior chemotherapy in the metastatic setting or developed disease recurrence during or within 6 months of completing adjuvant chemotherapy
- Unresectable or metastatic non-small cell lung cancer (NSCLC) whose tumors have activating HER2 (ERBB2) mutations, as detected by an FDA-approved test, and who have received a prior systemic therapy

This indication is approved under accelerated approval based on objective response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

- Locally advanced or metastatic HER2-positive (IHC 3+ or IHC 2+/ISH positive) gastric or gastroesophageal junction (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen
- Unresectable or metastatic HER2-positive (IHC 3+) solid tumors who have received prior systemic treatment and have no satisfactory alternative treatment options

This indication is approved under accelerated approval based on objective response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

WARNING: INTERSTITIAL LUNG DISEASE and EMBRYO-FETAL TOXICITY

- **Interstitial lung disease (ILD) and pneumonitis, including fatal cases, have been reported with ENHERTU. Monitor for and promptly investigate signs and symptoms including cough, dyspnea, fever, and other new or worsening respiratory symptoms. Permanently discontinue ENHERTU in all patients with Grade 2 or higher ILD/pneumonitis. Advise patients of the risk and to immediately report symptoms.**
- **Exposure to ENHERTU during pregnancy can cause embryo-fetal harm. Advise patients of these risks and the need for effective contraception.**

Contraindications

None.

Warnings and Precautions

Interstitial Lung Disease / Pneumonitis

Severe, life-threatening, or fatal interstitial lung disease (ILD), including pneumonitis, can occur in patients treated with ENHERTU. A higher incidence of Grade 1 and 2 ILD/pneumonitis has been observed in patients with moderate renal impairment. Advise patients to immediately report cough, dyspnea, fever, and/or any new or worsening respiratory symptoms. Monitor patients for signs and symptoms of ILD. Promptly investigate evidence of ILD. Evaluate patients with suspected ILD by radiographic imaging. Consider consultation with a pulmonologist. For asymptomatic ILD/pneumonitis (Grade 1), interrupt ENHERTU until resolved to Grade 0, then if resolved in ≤ 28 days from date of onset, maintain dose. If resolved in > 28 days from date of onset, reduce dose 1 level. Consider corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g., ≥ 0.5 mg/kg/day prednisolone or equivalent). For symptomatic ILD/pneumonitis (Grade 2 or greater), permanently discontinue ENHERTU. Promptly initiate systemic corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g., ≥ 1 mg/kg/day prednisolone or equivalent) and continue for at least 14 days followed by gradual taper for at least 4 weeks.

HER2-Positive, HER2-Low, and HER2-Ultralow Metastatic Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

In patients with metastatic breast cancer, HER2-mutant NSCLC, and other solid tumors treated with ENHERTU 5.4 mg/kg, ILD occurred in 12% of patients. Median time to first onset was 5.5 months (range: 0.9 to 31.5). Fatal outcomes due to ILD and/or pneumonitis occurred in 0.9% of patients treated with ENHERTU.

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, ILD occurred in 10% of patients. Median time to first onset was 2.8 months (range: 1.2 to 21).

Neutropenia

Severe neutropenia, including febrile neutropenia, can occur in patients treated with ENHERTU. Monitor

complete blood counts prior to initiation of ENHERTU and prior to each dose, and as clinically indicated. For Grade 3 neutropenia (Absolute Neutrophil Count [ANC] <1.0 to $0.5 \times 10^9/L$), interrupt ENHERTU until resolved to Grade 2 or less, then maintain dose. For Grade 4 neutropenia (ANC $<0.5 \times 10^9/L$), interrupt ENHERTU until resolved to Grade 2 or less, then reduce dose by 1 level. For febrile neutropenia (ANC $<1.0 \times 10^9/L$ and temperature $>38.3^\circ C$ or a sustained temperature of $\geq 38^\circ C$ for more than 1 hour), interrupt ENHERTU until resolved, then reduce dose by 1 level.

HER2-Positive, HER2-Low, and HER2-Ultralow Metastatic Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

In patients with metastatic breast cancer, HER2-mutant NSCLC, and other solid tumors treated with ENHERTU 5.4 mg/kg, a decrease in neutrophil count was reported in 65% of patients. Nineteen percent had Grade 3 or 4 decreased neutrophil count. Median time to first onset of decreased neutrophil count was 22 days (range: 2 to 939). Febrile neutropenia was reported in 1.2% of patients.

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, a decrease in neutrophil count was reported in 72% of patients. Fifty-one percent had Grade 3 or 4 decreased neutrophil count. Median time to first onset of decreased neutrophil count was 16 days (range: 4 to 187). Febrile neutropenia was reported in 4.8% of patients.

Left Ventricular Dysfunction

Patients treated with ENHERTU may be at increased risk of developing left ventricular dysfunction. Left ventricular ejection fraction (LVEF) decrease has been observed with anti-HER2 therapies, including ENHERTU. Assess LVEF prior to initiation of ENHERTU and at regular intervals during treatment as clinically indicated. Manage LVEF decrease through treatment interruption. When LVEF is $>45\%$ and absolute decrease from baseline is 10-20%, continue treatment with ENHERTU. When LVEF is 40-45% and absolute decrease from baseline is 20%, interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF of 20% is confirmed, permanently discontinue ENHERTU. Permanently discontinue ENHERTU in patients with symptomatic congestive heart failure. Treatment with ENHERTU has not been studied in patients with a history of clinically significant cardiac disease or LVEF $<50\%$ prior to initiation of treatment.

HER2-Positive, HER2-Low, and HER2-Ultralow Metastatic Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

In patients with metastatic breast cancer, HER2-mutant NSCLC, and other solid tumors treated with ENHERTU 5.4 mg/kg, LVEF decrease was reported in 4.6% of patients, of which 0.6% were Grade 3 or 4.

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, no clinical adverse events of heart failure were reported; however, on echocardiography, 8% were found to have asymptomatic Grade 2 decrease in LVEF.

Embryo-Fetal Toxicity

ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. Verify the pregnancy status of females of reproductive potential prior to the initiation of ENHERTU. Advise females of reproductive potential to use effective contraception during treatment and for 7 months after the last dose of ENHERTU. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for 4 months after the last dose of ENHERTU.

Additional Dose Modifications

Thrombocytopenia

For Grade 3 thrombocytopenia (platelets <50 to $25 \times 10^9/L$) interrupt ENHERTU until resolved to Grade 1 or less, then maintain dose. For Grade 4 thrombocytopenia (platelets $<25 \times 10^9/L$) interrupt ENHERTU until resolved to Grade 1 or less, then reduce dose by 1 level.

Adverse Reactions

HER2-Positive, HER2-Low, and HER2-Ultralow Metastatic Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

The pooled safety population reflects exposure to ENHERTU 5.4 mg/kg intravenously every 3 weeks in 2233 patients in Study DS8201-A-J101 (NCT02564900), DESTINY-Breast01, DESTINY-Breast02, DESTINY-Breast03, DESTINY-Breast04, DESTINY-Breast06, DESTINY-Lung01, DESTINY-Lung02, DESTINY-CRC02, and DESTINY-PanTumor02. Among these patients, 67% were exposed for >6 months and 38% were exposed for >1 year. In this pooled safety population, the most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (73%), nausea (72%), decreased hemoglobin (67%), decreased neutrophil count (65%), decreased lymphocyte count (60%), fatigue (55%), decreased platelet count (48%), increased aspartate aminotransferase (46%), increased alanine aminotransferase (44%), increased blood alkaline phosphatase (39%), vomiting (38%), alopecia (37%), constipation (32%), decreased blood potassium (32%), decreased appetite (31%), diarrhea (30%), and musculoskeletal pain (24%).

HER2-Positive Metastatic Breast Cancer

DESTINY-Breast03

The safety of ENHERTU was evaluated in 257 patients with unresectable or metastatic HER2-positive breast cancer who received at least 1 dose of ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast03. The median duration of treatment was 14 months (range: 0.7 to 30) for patients who received ENHERTU.

Serious adverse reactions occurred in 19% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were vomiting, ILD, pneumonia, pyrexia, and urinary tract infection. Fatalities due to adverse reactions occurred in 0.8% of patients including COVID-19 and sudden death (1 patient each).

ENHERTU was permanently discontinued in 14% of patients, of which ILD/pneumonitis accounted for 8%. Dose interruptions due to adverse reactions occurred in 44% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, leukopenia, anemia, thrombocytopenia, pneumonia, nausea, fatigue, and ILD/pneumonitis. Dose reductions occurred in 21% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were nausea, neutropenia, and fatigue.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (74%), decreased neutrophil count (70%), increased aspartate aminotransferase (67%), decreased hemoglobin (64%), decreased lymphocyte count (55%), increased alanine aminotransferase (53%), decreased platelet count (52%), fatigue (49%), vomiting (49%), increased blood alkaline phosphatase (49%), alopecia (37%), decreased blood potassium (35%), constipation (34%), musculoskeletal pain (31%), diarrhea (29%), decreased appetite (29%), headache (22%), respiratory infection (22%), abdominal pain (21%), increased blood bilirubin (20%), and stomatitis (20%).

HER2-Low and HER2-Ultralow Metastatic Breast Cancer

DESTINY-Breast06

The safety of ENHERTU was evaluated in 434 patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) or HER2-ultralow (IHC 0 with membrane staining) breast cancer who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast06. The median duration of treatment was 11 months (range: 0.4 to 39.6) for patients who received ENHERTU.

Serious adverse reactions occurred in 20% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, COVID-19, febrile neutropenia, and hypokalemia. Fatalities due to adverse reactions occurred in 2.8% of patients including ILD (0.7%); sepsis (0.5%); and COVID-19 pneumonia, bacterial meningoenzephalitis, neutropenic sepsis, peritonitis, cerebrovascular accident, general physical health deterioration (0.2% each).

ENHERTU was permanently discontinued in 14% of patients. The most frequent adverse reaction (>2%) associated with permanent discontinuation was ILD/pneumonitis. Dose interruptions due to adverse reactions occurred in 48% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were COVID-19, decreased neutrophil count, anemia, pyrexia, pneumonia, decreased white blood cell count, and ILD. Dose reductions occurred in 25% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were nausea, fatigue, decreased platelet count, and decreased neutrophil count.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (86%), decreased neutrophil count (75%), nausea (70%), decreased hemoglobin (69%), decreased lymphocyte count (66%), fatigue (53%), decreased platelet count (48%), alopecia (48%), increased alanine aminotransferase (44%), increased blood alkaline phosphatase (43%), increased aspartate aminotransferase (41%), decreased blood potassium (35%), diarrhea (34%), vomiting (34%), constipation (32%), decreased appetite (26%), COVID-19 (26%), and musculoskeletal pain (24%).

DESTINY-Breast04

The safety of ENHERTU was evaluated in 371 patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) breast cancer who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast04. The median duration of treatment was 8 months (range: 0.2 to 33) for patients who received ENHERTU.

Serious adverse reactions occurred in 28% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, pneumonia, dyspnea, musculoskeletal pain, sepsis, anemia, febrile neutropenia, hypercalcemia, nausea, pyrexia, and vomiting. Fatalities due to adverse reactions occurred in 4% of patients including ILD/pneumonitis (3 patients); sepsis (2 patients); and ischemic colitis, disseminated intravascular coagulation, dyspnea, febrile neutropenia, general physical health deterioration, pleural effusion, and respiratory failure (1 patient each).

ENHERTU was permanently discontinued in 16% of patients, of which ILD/pneumonitis accounted for 8%. Dose interruptions due to adverse reactions occurred in 39% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, fatigue, anemia, leukopenia, COVID-19, ILD/pneumonitis, increased transaminases, and hyperbilirubinemia. Dose reductions occurred in 23% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were fatigue, nausea, thrombocytopenia, and neutropenia.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (70%), decreased hemoglobin (64%), decreased neutrophil count (64%), decreased lymphocyte count (55%), fatigue (54%), decreased platelet count (44%), alopecia (40%), vomiting (40%), increased aspartate aminotransferase (38%), increased alanine aminotransferase (36%), constipation (34%), increased blood alkaline phosphatase (34%), decreased appetite (32%), musculoskeletal pain (32%), diarrhea (27%), and decreased blood potassium (25%).

HER2-Mutant Unresectable or Metastatic NSCLC (5.4 mg/kg)

DESTINY-Lung02 evaluated 2 dose levels (5.4 mg/kg [n=101] and 6.4 mg/kg [n=50]); however, only the results for the recommended dose of 5.4 mg/kg intravenously every 3 weeks are described below due to increased toxicity observed with the higher dose in patients with NSCLC, including ILD/pneumonitis.

The safety of ENHERTU was evaluated in 101 patients with HER2-mutant unresectable or metastatic NSCLC who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks until disease progression or unacceptable toxicity in DESTINY-Lung02. Nineteen percent of patients were exposed for >6 months.

Serious adverse reactions occurred in 30% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, thrombocytopenia, dyspnea, nausea, pleural effusion, and increased troponin I. Fatality occurred in 1 patient with suspected ILD/pneumonitis (1%).

ENHERTU was permanently discontinued in 8% of patients. Adverse reactions which resulted in permanent discontinuation of ENHERTU were ILD/pneumonitis, diarrhea, decreased blood potassium, hypomagnesemia, myocarditis, and vomiting. Dose interruptions of ENHERTU due to adverse reactions occurred in 23% of patients. Adverse reactions which required dose interruption (>2%) included neutropenia and ILD/pneumonitis. Dose reductions due to an adverse reaction occurred in 11% of patients.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were nausea (61%), decreased white blood cell count (60%), decreased hemoglobin (58%), decreased neutrophil count (52%), decreased lymphocyte count (43%), decreased platelet count (40%), decreased albumin (39%), increased aspartate aminotransferase (35%), increased alanine aminotransferase (34%), fatigue (32%), constipation (31%), decreased appetite (30%), vomiting (26%), increased alkaline phosphatase (22%), and alopecia (21%).

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

The safety of ENHERTU was evaluated in 187 patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma in DESTINY-Gastric01. Patients intravenously received at least 1 dose of either ENHERTU (N=125) 6.4 mg/kg every 3 weeks or either irinotecan (N=55) 150 mg/m² biweekly or paclitaxel (N=7) 80 mg/m² weekly for 3 weeks. The median duration of treatment was 4.6 months (range: 0.7 to 22.3) for patients who received ENHERTU.

Serious adverse reactions occurred in 44% of patients receiving ENHERTU 6.4 mg/kg. Serious adverse reactions in >2% of patients who received ENHERTU were decreased appetite, ILD, anemia, dehydration, pneumonia, cholestatic jaundice, pyrexia, and tumor hemorrhage. Fatalities due to adverse reactions occurred in 2.4% of patients: disseminated intravascular coagulation, large intestine perforation, and pneumonia occurred in 1 patient each (0.8%).

ENHERTU was permanently discontinued in 15% of patients, of which ILD accounted for 6%. Dose interruptions due to adverse reactions occurred in 62% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, anemia, decreased appetite, leukopenia, fatigue, thrombocytopenia, ILD, pneumonia, lymphopenia, upper respiratory tract infection, diarrhea, and decreased blood potassium. Dose reductions occurred in 32% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were neutropenia, decreased appetite, fatigue, nausea, and febrile neutropenia.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased hemoglobin (75%), decreased white blood cell count (74%), decreased neutrophil count (72%), decreased lymphocyte count (70%), decreased platelet count (68%), nausea (63%), decreased appetite (60%), increased aspartate aminotransferase (58%), fatigue (55%), increased blood alkaline phosphatase (54%), increased alanine aminotransferase (47%), diarrhea (32%), decreased blood potassium (30%), vomiting (26%), constipation (24%), increased blood bilirubin (24%), pyrexia (24%), and alopecia (22%).

HER2-Positive (IHC 3+) Unresectable or Metastatic Solid Tumors

The safety of ENHERTU was evaluated in 347 adult patients with unresectable or metastatic HER2-positive (IHC 3+) solid tumors who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast01, DESTINY-PanTumor02, DESTINY-Lung01, and DESTINY-CRC02. The median duration of treatment was 8.3 months (range 0.7 to 30.2).

Serious adverse reactions occurred in 34% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were sepsis, pneumonia, vomiting, urinary tract infection, abdominal pain, nausea, pneumonitis, pleural effusion, hemorrhage, COVID-19, fatigue, acute kidney injury, anemia, cellulitis, and dyspnea. Fatalities due to adverse reactions occurred in 6.3% of patients including ILD/pneumonitis (2.3%), cardiac arrest (0.6%), COVID-19 (0.6%), and sepsis (0.6%). The following events occurred in 1 patient each (0.3%): acute kidney injury, cerebrovascular accident, general physical health deterioration, pneumonia, and hemorrhagic shock.

ENHERTU was permanently discontinued in 15% of patients, of which ILD/pneumonitis accounted for 10%. Dose interruptions due to adverse reactions occurred in 48% of patients. The most frequent adverse reactions (>2%) associated with dose interruption were decreased neutrophil count, anemia, COVID-19, fatigue, decreased white blood cell count, and ILD/pneumonitis. Dose reductions occurred in 27% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were fatigue, nausea, decreased neutrophil count, ILD/pneumonitis, and diarrhea.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (75%), nausea (69%), decreased hemoglobin (67%), decreased neutrophil count (66%), fatigue (59%), decreased lymphocyte count (58%), decreased platelet count (51%), increased aspartate aminotransferase (45%), increased alanine aminotransferase (44%), increased blood alkaline phosphatase (36%), vomiting (35%), decreased appetite (34%), alopecia (34%), diarrhea (31%), decreased blood potassium (29%), constipation (28%), decreased sodium (22%), stomatitis (20%), and upper respiratory tract infection (20%).

Use in Specific Populations

- **Pregnancy:** ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. There are clinical considerations if ENHERTU is used in pregnant women, or if a patient becomes pregnant within 7 months after the last dose of ENHERTU.
- **Lactation:** There are no data regarding the presence of ENHERTU in human milk, the effects on the breastfed child, or the effects on milk production. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with ENHERTU and for 7 months after the last dose.
- **Females and Males of Reproductive Potential:** Pregnancy testing: Verify pregnancy status of females of reproductive potential prior to initiation of ENHERTU. Contraception: *Females:* ENHERTU can cause fetal harm when administered to a pregnant woman. Advise females of reproductive potential to use effective contraception during treatment with ENHERTU and for 7 months after the last dose. *Males:* Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for 4 months after the last dose. Infertility: ENHERTU may impair male reproductive function and fertility.
- **Pediatric Use:** Safety and effectiveness of ENHERTU have not been established in pediatric patients.
- **Geriatric Use:** Of the 1741 patients with HER2-positive, HER2-low, or HER2-ultralow breast cancer treated with ENHERTU 5.4 mg/kg, 24% were ≥ 65 years and 4.9% were ≥ 75 years. No overall differences in efficacy within clinical studies were observed between patients ≥ 65 years of age compared to younger patients. There was a higher incidence of Grade 3-4 adverse reactions observed in patients aged ≥ 65 years (61%) as compared to younger patients (52%). Of the 101 patients with HER2-mutant unresectable or metastatic NSCLC treated with ENHERTU 5.4 mg/kg, 40% were ≥ 65 years and 8% were ≥ 75 years. No overall differences in efficacy or safety were observed between patients ≥ 65 years of age compared to younger patients. Of the 125 patients with HER2-positive locally advanced or metastatic gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg in DESTINY-Gastric01, 56% were ≥ 65 years and 14% were ≥ 75 years. No overall differences in efficacy or safety were observed between patients ≥ 65 years of age compared to younger patients. Of the 192 patients with HER2-positive (IHC 3+) unresectable or metastatic solid tumors treated with ENHERTU 5.4 mg/kg in DESTINY-PanTumor02, DESTINY-Lung01, or DESTINY-CRC02, 39% were ≥ 65 years and 9% were ≥ 75 years. No overall differences in efficacy or safety were observed between patients ≥ 65 years of age compared to younger patients.
- **Renal Impairment:** A higher incidence of Grade 1 and 2 ILD/pneumonitis has been observed in patients with moderate renal impairment. Monitor patients with moderate renal impairment more frequently. The recommended dosage of ENHERTU has not been established for patients with severe renal impairment ($CL_{Cr} < 30$ mL/min).
- **Hepatic Impairment:** In patients with moderate hepatic impairment, due to potentially increased exposure, closely monitor for increased toxicities related to the topoisomerase inhibitor, DXd. The recommended dosage of ENHERTU has not been established for patients with severe hepatic

impairment (total bilirubin >3 times ULN and any AST).

To report SUSPECTED ADVERSE REACTIONS, contact Daiichi Sankyo, Inc. at 1-877-437-7763 or FDA at 1-800-FDA-1088 or [fda.gov/medwatch](https://www.fda.gov/medwatch).

Please see accompanying full [Prescribing Information](#), including Boxed WARNINGS, and [Medication Guide](#).

About Daiichi Sankyo

Daiichi Sankyo is an innovative global healthcare company contributing to the sustainable development of society that discovers, develops and delivers new standards of care to enrich the quality of life around the world. With more than 120 years of experience, Daiichi Sankyo leverages its world-class science and technology to create new modalities and innovative medicines for people with cancer, cardiovascular and other diseases with high unmet medical need. For more information, please visit www.daiichisankyo.com.

Media Contacts:

Global/US:

Jennifer Brennan
Daiichi Sankyo, Inc.
jennifer.brennan@daiichisankyo.com
+1 908 900 3183 (mobile)

Japan:

Daiichi Sankyo Co., Ltd.
DS-PR_jp@daiichisankyo.com

Investor Relations Contact:

DaiichiSankyoIR_jp@daiichisankyo.com

References:

-
- ¹ Bray F, et al. *CA Cancer J Clin*. 2024; 10.3322/caac.21834.
 - ² American Cancer Society. *Key Statistics for Breast Cancer*. Accessed January 2025.
 - ³ National Cancer Institute. *SEER Cancer Stat Facts: Female Breast Cancer Subtypes*. Accessed January 2025.
 - ⁴ Iqbal N, et al. *Mol Biol Int*. 2014;852748.
 - ⁵ Ahn S, et al. *J Pathol Transl Med*. 2020;54(1):34-44.
 - ⁶ Sajjadi E, et al. *Cancer Drug Resist*. 2022;5(4):882-888.
 - ⁷ Manohar P, et al. *Cancer Biol Med*. 2022 Feb 15; 19(2):202–212.
 - ⁸ Cortes J, et al. *Lancet*. 2011;377:914-923.
 - ⁹ Yuan P, et al. *Eur J Cancer*. 2019;112:57-65.
 - ¹⁰ Jerusalem G, et al. *JAMA Oncol*. 2018;4(10):1367–1374.
 - ¹¹ Modi S, et al. *N Engl J Med*. 2022;387:9-20.
 - ¹² Eiger D, et al. *Cancers*. 2021 Mar; 13(5): 1015.